

# Whole Exome Sequencing (WES)

Procedure(s) addressed by this policy:	Procedure Code(s)
Exome (e.g., unexplained constitutional or heritable disorder or syndrome); sequence analysis	81415
Sequence analysis, each comparator exome (e.g., parent(s), sibling(s))	81416
Re-evaluation of previously obtained exome sequence (e.g., updated knowledge or unrelated condition/syndrome)	81417

## What Is Whole Exome Sequencing?

- Whole exome sequencing (WES) utilizes DNA-enrichment methods and massively parallel nucleotide sequencing to identify disease-associated variants throughout the human genome.
- WES has been proposed for diagnostic use in individuals who present with complex genetic phenotypes suspected of having a rare genetic condition, who cannot be diagnosed by standard clinical workup, or when features suggest a broad differential diagnosis that would require evaluation by multiple genetic tests.
- The standard approach to the diagnostic evaluation of an individual suspected of having a rare genetic condition may include combinations of radiographic, biochemical, electrophysiologic, and targeted genetic testing such as a chromosomal microarray, single-gene analysis, and/or a targeted gene panel.<sup>1</sup>
- WES is typically not an appropriate first-tier test, but can be appropriate if initial testing is unrevealing, or if there is no single-gene or panel test available for the particular condition.<sup>2</sup>
- Identifying a molecularly confirmed diagnosis in a timely manner for an individual with a rare genetic condition can have a variety of health outcomes<sup>2,3,4,5,6,7,8,9</sup>, including:
  - guiding prognosis and improving clinical decision-making, which can improve clinical outcome by
    - application of specific treatments as well as withholding of contraindicated treatments for certain rare genetic conditions
    - surveillance for later-onset comorbidities
    - initiation of palliative care
    - withdrawal of care
  - reducing the financial & psychological impact of diagnostic uncertainty and the diagnostic odyssey (e.g., eliminating lower-yield testing and additional screening testing that may later be proven unnecessary once a diagnosis is achieved)
  - informing genetic counseling related to recurrence risk and prenatal diagnosis options



- allowing for more rapid molecular diagnosis than a sequential genetic testing approach

## Test Information

- WES is limited to the DNA sequence of coding regions (exons) and flanking intronic regions of the genome, which is estimated to contain 85% of heritable disease-causing variants.
- Pathogenic variants that can be identified by WES include missense, nonsense, splice-site, and small deletions or insertions.
- At the present time, WES will typically miss certain classes of disease-causing variants, such as structural variants (e.g., translocations, inversions), abnormal chromosome imprinting or methylation, copy-number variants, some mid-size insertions and deletions (ca. 10-500 bp), trinucleotide repeat expansion mutations, deeper intronic mutations, and low-level mosaicism.
- WES has the advantage of decreased turnaround time and increased efficiency relative to Sanger sequencing of multiple genes.
- WES is associated with technical and analytical variability, including uneven sequencing coverage, gaps in exon capture before sequencing, as well as variability in variant classification based on proprietary filtering algorithms and potential lack of critical clinical history or family samples.<sup>10</sup>

## Guidelines and Evidence

- The American College of Medical Genetics and Genomics, Association for Molecular Pathology, and College of American Pathologists convened a workgroup that developed standard terminology for describing sequence variants. The guidelines describe criteria for classifying sequence variants into five categories (pathogenic, likely pathogenic, uncertain significance, likely benign, and benign) based on criteria using typical types of variant evidence (e.g., population data, computational data, functional data, segregation data).<sup>11</sup>
- The American College of Medical Genetics has three relevant policy statements that offer guidance on: 1) the clinical application of whole exome and whole genome testing,<sup>12</sup> 2) informed consent for genome/exome sequencing,<sup>13</sup> and 3) reporting of incidental findings in clinical exome and genome sequencing.<sup>14,15</sup>
- Evidence for the clinical utility of WES in individuals with multiple congenital anomalies and/or a neurodevelopmental phenotype includes numerous large case series. Relevant outcomes include improved clinical decision-making (e.g., application of specific treatments, withholding of contraindicated treatments, changes to surveillance), changes in reproductive decision making, and resource utilization. WES serves as a powerful diagnostic tool for individuals with rare genetic conditions in which the specific genetic etiology is unclear or unidentified by standard clinical workup.<sup>7,16,17</sup>



- The average diagnostic yield of WES is 20-40% depending on the individual's age, phenotype, previous workup, and number of comparator samples analyzed.<sup>5,16,18</sup> Among individuals with a pathogenic or likely pathogenic findings by WES, 5-7% received a dual molecular diagnosis (i.e., two significant findings associated with non-overlapping clinical presentations).<sup>16,18</sup>
- The use of family trio WES reduces the rate of uncertain findings, adds to the clinical sensitivity with regard to the interpretation of clinically novel genes, and increases the diagnostic utility of WES. For example, in three publications the positive rate ranges from 31-37% in patients undergoing trio analysis compared to 20-23% positive rate among proband-only WES.<sup>16,19,20</sup>
- Re-evaluation of previously obtained exome sequence has the potential for additional diagnostic yield because of constant expansions of existing variant databases, as well as periodic novel gene discovery and publication.<sup>21</sup>

## Criteria

- Whole exome sequencing (WES) is considered medically necessary for the evaluation of unexplained congenital or neurodevelopmental disorder in children <21 years of age when ALL of the following criteria are met:
  - The patient and family history have been evaluated by a Board-Certified or Board-Eligible Medical Geneticist, or an Advanced Practice Nurse in Genetics (APGN) credentialed by either the Genetic Nursing Credentialing Commission (GNCC) or the American Nurses Credentialing Center (ANCC), and
  - A genetic etiology is considered the most likely explanation for the phenotype, based on EITHER of the following, and
    - multiple congenital abnormalities affecting unrelated organ systems
    - TWO of the following criteria are met:
      - abnormality affecting at minimum a single organ system
      - significant developmental delay, intellectual disability (e.g., characterized by significant limitations in both intellectual functioning and in adaptive behavior), symptoms of a complex neurodevelopmental disorder (e.g., self-injurious behavior, reverse sleep-wake cycles, dystonia, hemiplegia, spasticity, epilepsy, muscular dystrophy), and/or severe neuropsychiatric condition (e.g., schizophrenia, bipolar disorder, Tourette syndrome)
      - family history strongly suggestive of a genetic etiology, including consanguinity
      - period of unexplained developmental regression
      - biochemical findings suggestive of an inborn error of metabolism
  - Alternate etiologies have been considered and ruled out when possible (e.g., environmental exposure, injury, infection), and



- Clinical presentation does not fit a well-described syndrome for which single-gene or targeted panel testing (e.g., comparative genomic hybridization [CGH]/chromosomal microarray analysis [CMA]) is available, and
- WES is more efficient and economical than the separate single-gene tests or panels that would be recommended based on the differential diagnosis (e.g., genetic conditions that demonstrate a high degree of genetic heterogeneity), and
- A diagnosis cannot be made by standard clinical work-up, excluding invasive procedures such as muscle biopsy, and
- Predicted impact on health outcomes, as above, and
- Pre- and post-test counseling by an appropriate provider (as deemed by the Health Plan policy), such as an American Board of Medical Genetics or American Board of Genetic Counseling-certified Genetic Counselor

### Exclusions and Other Considerations:

- WES is considered experimental/investigational for the diagnosis of genetic disorders in individuals <21 years of age who do not meet the above criteria.
- WES is considered experimental/investigational for screening for genetic disorders in asymptomatic or pre-symptomatic individuals.
- Ideal sample type should be considered based on the clinical presentation (e.g., suspect mosaicism based on pigmentary anomalies, consider skin fibroblast as ideal sample type).

### References

1. Shashi V, McConkie-Rosell A, Rosell B, et al. The utility of the traditional medical genetics diagnostic evaluation in the context of next-generation sequencing for undiagnosed genetic disorders. *Genet Med.* 2014 Feb; 16: 176–182. PMID: 23928913
2. Sawyer SL, Hartley T, Dymant DA et al. Utility of whole-exome sequencing for those near the end of the diagnostic odyssey: time to address gaps in care. *Clin Genet.* 2016 Mar; 89: 275–284. PMID:26283276
3. Dixon-Salazar TJ, Silhavy JL, Udpa N, et al. Exome sequencing can improve diagnosis and alter patient management. *Sci Transl Med.* Jun 13 2012;4(138):138ra178. PMID 22700954
4. Halverson CM, Clift KE, & JB McCormick. Was it worth it? Patients’ perspectives on the perceived value of genomic-based individualized medicine. *J Community Genet.* Apr 2016; 7(2):145–152. PMID 26860291
5. Iglesias A, Anyane-Yeboah K, Wynn J, et al. The usefulness of whole-exome sequencing in routine clinical practice. *Genet Med.* Dec 2014;16(12):922-931. PMID24901346
6. Nolan D, Carlson M. Whole exome sequencing in pediatric neurology patients: clinical implications and estimated cost analysis. *J Child Neurol.* Jun 2016;31(7):887-894. PMID 26863999
7. Stark Z, Tan TY, Chong B, et al. A prospective evaluation of whole-exome sequencing as a first-tier molecular test in infants with suspected monogenic disorders. *Genet Med.* Nov 2016;18(11):1090-1096. PMID26938784



8. Thevenon J, Duffourd Y, Masurel-Paulet A, et al. Diagnostic odyssey in severe neurodevelopmental disorders: toward clinical whole-exome sequencing as a first-line diagnostic test. *Clinical Genetics*. 2016 Jun; 89: 700–707. PMID: 26757139.
9. Valencia CA, Husami A, Holle J, et al. Clinical impact and cost-effectiveness of whole exome sequencing as a diagnostic tool: a pediatric center's experience. *Front Pediatr*. 2015;3:67. PMID 26284228
10. Rehm HL, Bale SJ, Bayrak-Toydemir P, et al. ACMG clinical laboratory standards for next-generation sequencing. *Genet Med*. Sep 2013;15(9):733-747. PMID 23887774
11. Richards S, Aziz N, Bale S, et al. Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology. *Genet Med*. May 2015;17(5):405-424. PMID 25741868
12. ACMG Board of Directors. Points to consider in the clinical application of whole-genome sequencing. *Genet Med* 2012; 14:759–761. PMID 22863877
13. ACMG Board of Directors. Points to consider for informed consent for genome/exome sequencing. *Genet Med* Sept 2013;15(9):748–749. PMID 23970068
14. Green RC, Berg JS, Grody WW, et al. ACMG recommendations for reporting of incidental findings in clinical exome and genome sequencing. *Genet Med*. Jul 2013;15(7):565-574. PMID 23788249
15. Kalia SS, Adelman K, Bale SJ, et al. Recommendations for reporting of secondary findings in clinical exome and genome sequencing, 2016 update (ACMG SF v2.0): a policy statement of the American College of Medical Genetics and Genomics. *Genet Med*. 2016 Nov 17. doi: 10.1038/gim.2016.190. [Epub ahead of print] PMID: 27854360
16. Farwell KD, Shahmirzadi L, El-Khechen D, et al. Enhanced utility of family-centered diagnostic exome sequencing with inheritance model-based analysis: results from 500 unselected families with undiagnosed genetic conditions. *Genet Med*. Jul 2015;17(7):578-586. PMID 25356970
17. Yang Y, Muzny DM, Xia F, et al. Molecular Findings Among Patients Referred for Clinical Whole-Exome Sequencing. *JAMA*. 2014;312(18):1870-1879. PMID 25326635
18. Trujillano D, Bertoli-Avella AM, Kumar Kandaswamy K, et al. Clinical exome sequencing: results from 2819 samples reflecting 1000 families. *Eur J Hum Genet*. 2017 Feb;25(2):176-182. PMID 27848944
19. Lee H, Deignan JL, Dorrani N, et al. Clinical exome sequencing for genetic identification of rare Mendelian disorders. *JAMA*. Nov 12 2014;312(18):1880-1887. PMID 25326637
20. Retterer K, Jussola J, Cho MT, et al. Clinical application of whole-exome sequencing across clinical indications. *Genet Med*. Dec 2016; 18: 696–704. PMID 26633542
21. Wenger AM, Guturu H, Bernstein JA, et al. Systematic reanalysis of clinical exome data yields additional diagnoses: implications for providers. *Genet med*. Jul 2016. Epub, ahead of print. PMID 27441994

