

Inherited Bone Marrow Failure (BMF) Syndromes Panel Testing

Procedure(s) addressed by this policy:	Procedure Code(s)
Unlisted molecular pathology procedure	81479

What are inherited bone marrow failure syndromes?

Bone marrow failure (BMF) is the inability of the bone marrow to produce sufficient numbers of functional blood cells to meet physiologic demands and is typically classified into three categories, based on presumed etiology, including inherited, secondary, or idiopathic.¹ Inherited bone marrow failure syndromes (IBMFSs) are a group of genetically defined disorders that are characterized by BMF. IBMFSs typically present with cytopenias and are often associated with other congenital anomalies and/or physical features.¹ Patients presenting with aplastic anemia (AA), myelodysplastic syndrome (MDS), acute myeloid leukemia (AML), and chronic unexplained cytopenias should be evaluated for an IBMFS.¹

- While specific features vary by each type of IBMFS, features that are present in most IBMFSs include bone marrow failure with single or multi-lineage cytopenia and many carry an increased risk to develop aplastic anemia (AA), myelodysplastic syndrome (MDS), acute myeloid leukemia (AML), and solid malignancies.^{1,2}
- IBMFSs typically present within the first decade of life; however, delay in diagnosis and variability in phenotypic spectrum may lead to diagnosis even into adulthood.²
- IBMFSs have been shown to present with isolated aplastic anemia or myelodysplastic syndrome. An underlying IBMFS diagnosis was identified in 5.1% and 13.6% of presumed sporadic aplastic anemia and MDS cases, respectively.³
- Phenotypic overlap between IBMFSs makes it difficult to establish a diagnosis based solely on clinical features.²
- Treatment of IBMFSs varies depending on the specific type, but typically involves supportive care, including blood and/or specific blood cell transfusions, and in severe situations, hematopoietic stem cell transplants (HSCTs).
- Clinical management and outcomes in individuals with aplastic anemia and myelodysplastic syndrome differ in those with an underlying IBMFS compared to sporadic disease.¹
- Timely genetic testing is essential to establish a diagnosis in the proband and to guide appropriate management, treatment, and cancer surveillance.² Additionally, knowing the genetic cause in the proband allows for genetic testing



Seattle Children's[®]
HOSPITAL • RESEARCH • FOUNDATION

PLUGS[®]
Patient-centered Laboratory
Utilization Guidance Services

© 2012–2020 Seattle Children's Hospital. All rights reserved

June 2020

in family members. This information is important for their own health and a critical part of their workup if being considered as a possible bone marrow transplant donor.

- Some of the most common IBMFSs are summarized below:
 - **Fanconi Anemia (FA)** – A disorder that is typically inherited in an autosomal recessive manner, though X-linked and autosomal dominant forms have been reported. Heterozygotes (carriers) are not at risk for FA. However, carriers of a subset of FA-related genes (e.g., *BRCA2*, *PALB2*, and *BRIP1*) have an increased risk for breast and other cancers.⁴ FA is characterized by BMF and increased risk for malignancy. Physical abnormalities are present in ~75% of individuals and include growth deficiency, abnormal skin pigmentation, skeletal malformations of the upper and lower limbs (especially thumbs), microcephaly, ophthalmic anomalies, and genitourinary tract anomalies. Progressive BMF with pancytopenia is typically present in the first decade of life. Additional solid tumor malignancies include squamous cell cancers of the head, neck, and anogenital region. Diagnosis of FA is established in an individual with increased chromosome breakage and radial forms induced by diepoxybutane (DEB) and mitomycin C. Molecular testing can confirm the diagnosis with identification of pathogenic variant(s).⁴
 - **Telomere biology disorders** –Dyskeratosis congenita (DC) is the most common telomere biology disorder and is inherited in X-linked, autosomal dominant, or autosomal recessive manners. DC is caused by defective telomere maintenance and characterized by a classic triad of nail dysplasia, lacy reticular pigmentation of the upper chest/and or back, and oral leukoplakia, though this triad may not present until adolescence and adulthood.⁵ Infants may be severely affected with history of intrauterine growth restriction, cerebellar hypoplasia, immunodeficiency and retinopathy⁶. Another telomere biology disorder is caused by mutations in *TERC* and *TERT* and follows an autosomal dominant inheritance. Other physical features may include eye abnormalities, dental abnormalities, developmental delay, short stature, microcephaly, and genitourinary anomalies. It is recognized that there is a broad phenotypic spectrum ranging from mild to severe. Individuals with a telomere biology disorder are at increased risk for BMF, MDS, AML, solid tumors, pulmonary fibrosis, and liver cirrhosis. Pulmonary fibrosis is the most common presentation of a telomere biology disorder, and may be the only symptom in adults.⁷ Approximately 70% of individuals with a clinical diagnosis are found to have a pathogenic variant(s) in an associated gene.⁵
 - **Shwachman-diamond syndrome (SDS)** – SDS is a disorder most commonly inherited in an autosomal recessive fashion though some autosomal dominant cases have been reported. SDS is characterized by exocrine pancreatic dysfunction with gastrointestinal malabsorption,



malnutrition and growth failure. Individuals may also experience single or multi-lineage cytopenias and are at an increased risk for MDS and AML. A broad phenotypic spectrum has been recognized. Myers *et al.* found that only 19/37 patients with genetically confirmed SDS presented with the classical combination of features.⁸ Diagnosis can be established in an individual with the classic findings of exocrine pancreatic dysfunction and bone marrow dysfunction and/or by identification of pathogenic variant(s) in the *SBDS*, *ELF1*, *DNAJC21* or *SRP54* genes.⁹

- **Diamond-Blackfan anemia (DBA)** – DBA is a disorder typically inherited in an autosomal dominant manner, though *GATA1*- and *TSR2*-related DBA are inherited in an X-linked manner.¹⁰ Classical DBA is characterized by profound normochromic and typically macrocytic anemia. Approximately 25-50% of individuals with DBA will have congenital malformations including thumb malformations, craniofacial anomalies, and congenital heart disease; 30% will have growth retardation. Ninety percent of affected individuals will experience red cell aplasia within the first year of life. Other individuals have very mild anemia, requiring no treatment¹¹. There is an increased risk to develop AML, MDS, and solid tumors such as osteosarcoma. DBA is suspected in individuals who meet the following diagnostic criteria:
 1. Age younger than one year
 2. Macrocytic anemia with no other significant cytopenias
 3. Reticulocytopenia
 4. Normal marrow cellularity with a paucity of erythroid precursorsElevated erythrocyte adenosine deaminase activity (eADA) levels are elevated in the majority of individuals with DBA.¹¹ Genetic analysis of the 21 genes known to be associated with DBA will identify a pathogenic variant in ~65% of affected individuals.¹⁰
- **Severe congenital neutropenia (SCN)** – SCN is a “chronic state of severe neutropenia associated with a neutrophil count less than 500/μL lasting longer than 3 months, often presenting in the first year of life.”¹ SCN’s can be associated with different features, which typically include severe/recurrent infections, dental issues (chronic gingivitis and caries), decreased bone mineral density, and increased risk for MDS/AML. The most well-described SCN disorders are caused by pathogenic variants in the *HAX1* and *ELANE* genes. Additional pathogenic variants have been reported in *AK2*, *GF11*, *CSF3R*, *WAS*, and *G6PC3*. Autosomal dominant, autosomal recessive, and X-linked forms have been described.^{1,12}
- **Congenital amegakaryocytic thrombocytopenia (CAMT)** – CAMT is a rare autosomal recessive IBMFS associated with pathogenic variants in *MPL*. CAMT is characterized by isolated thrombocytopenia due to ineffective megakaryocytopoiesis at birth, with elevated plasma TPO levels.¹³ Thrombocytopenia will progress to pancytopenia/aplastic anemia



in the majority of affected individuals. Individuals are at risk to develop MDS and AML.^{13,14} Genotype-phenotype correlations exist, and individuals with type I variants have earlier progression to bone marrow failure than those with type II.¹³

- **GATA2 deficiency** – GATA2 deficiency is an autosomal dominant disorder caused by pathogenic variants in the *GATA2* gene. GATA2 deficiency results in a range of phenotypes including viral and bacterial infections, cytopenias, myelodysplasia, pulmonary alveolar proteinosis and lymphedema. Individuals with GATA2 deficiency have an increased risk to develop MDS and leukemias (AML and CMML).¹⁵ The majority of cases of pediatric patients who develop MDS will have a monosomy 7 bone marrow karyotype.¹⁶
- **SAMD9-related MIRAGE and SAMD9L-ATXPC syndromes** – Variants in the *SAMD9* and *SAMD9L* genes cause newly described IBMFSs, inherited in an autosomal dominant manner. *SAMD9* mutations cause MIRAGE (myelodysplasia, infection, restriction of growth, adrenal hyperplasia, genital phenotypes, and enteropathy) syndrome.¹⁷ *SAMD9L* mutations are characterized by cerebellar ataxia, cytopenia and predisposition to marrow failure with risk of MDS. Both syndromes are associated with monosomy 7 cytogenetic finding on bone marrow biopsy. These syndromes are likely underdiagnosed due to a common occurrence of genetic reversion to restore hematopoiesis.¹⁸
- **Familial myelodysplastic syndrome** – Myelodysplastic syndromes (MDS) are a heterogeneous group of disorders characterized by dysplastic changes in the bone marrow, cytopenias, and an increased risk to develop AML. MDS is primarily a sporadic disease that occurs in older individuals, but inherited forms have been described including the IBMFSs described above, as well as familial MDS syndromes associated with the *RUNX1*, *CEBPA*, *DDX41*, *ANKRD26*, and *ETV6* genes. Familial MDS disorders are inherited in an autosomal dominant manner.^{19,20}

Test Information

- The investigation and diagnosis of patients with IBMFSs necessitates a combination of laboratory analyses, including completed blood counts with differential, telomere length studies, exocrine pancreatic function studies, bone marrow analysis, and cytogenetic studies, along with clinical assessment and genetic testing.¹
- Clinical genetic testing is available for many IBMFSs, via single gene analysis and/or multi-gene panels. The efficiency of next generation sequencing (NGS) has led to an increasing number of large, multi-gene testing panels. NGS panels that test several genes at once are particularly well-suited to conditions caused by more than one gene or where there is considerable clinical overlap between conditions making it difficult to reliably narrow down likely causes. As a result, several laboratories have



Seattle Children's[®]
HOSPITAL • RESEARCH • FOUNDATION

PLUGS[®]

Patient-centered Laboratory
Utilization Guidance Services

© 2012–2020 Seattle Children's Hospital. All rights reserved

June 2020

begun to combine genes involved in certain conditions which often have both of those characteristics.

- Under certain circumstances, technologies used in multi-gene testing may fail to identify mutations that might be identifiable through single-gene testing. If high clinical suspicion remains for a particular syndrome after negative multi-gene test results, consultation with the testing lab and/or additional targeted genetic testing may be warranted.
- Multi-gene tests vary in technical specifications (e.g., depth of coverage, extent of intron/exon boundary analysis, methodology of large deletion/duplication analysis).

Guidelines and Evidence

- No current U.S. guidelines address the use of multi-gene panels in IBMFSs.
- An expert-authored review (2017) states the following regarding IBMFSs¹:
 - “Genetic testing is an indispensable tool in the diagnostic evaluation of IBMFSs that complements traditional clinical history, examination, and laboratory evaluation, especially in the setting of overlapping or adult presentations. However, clinical use of this powerful tool is currently limited by cost or access in most places.”
 - "In addition, even when genetic testing is available, it may fail to provide the correct diagnosis." This is because not all genes that cause IBMFS have been identified, many rare variants in known IBMFS genes cannot currently be classified as disease causing, or in the event of somatic reversion, the genetic variant(s) that cause a patient's IBMFS may not be detectable in peripheral blood cells.
 - "Now and likely well into the future, the sum of all available tools is greater than any alone, and a modern IBMFS workup should include a focused history and physical examination, screening tests, and genetic evaluation whenever possible."
- There are guidelines published for a subset of IBMFSs that have suggestive laboratory findings in addition to genetic sequencing:
 - Fanconi Anemia
 - The Fanconi Anemia Research Fund Inc. held a consensus conference in 2013 in which experts established guidelines for diagnosis and management of Fanconi anemia:
 - “Patients with FA may present with AA, MDS, single cytopenias, or macrocytic red cells without another explanation. A diagnosis of FA should be considered in all children and young adults with androgen-responsive or ATG/cyclosporine A-non-responsive “acquired” aplastic

anemia. It is absolutely imperative to test for FA if a stem cell transplant is planned, as standard stem cell transplant conditioning will result in significant morbidity and high mortality rates for patients with FA.”²¹

- “If the results from the chromosome breakage test are positive, then mutation analysis should be performed to identify the specific genetic mutation that has caused the patient to develop FA.”²¹
- “The first test that should be used to diagnose FA is the chromosome breakage test, which is performed on a sample of the patient’s blood in a clinical cytogenetics laboratory.”²¹ Recommendations for follow-up testing are made based on the results of the chromosome breakage studies:
 - Negative:
 - No further testing for Fanconi anemia unless strong clinical suspicion
 - Positive:
 - Next-generation sequencing for known FA genes
 - Copy number evaluation for deletions/duplications if not captured on original panel and no mutations are identified in gene panel
 - Equivocal:
 - Next-generation sequencing or targeted mutation testing for other chromosome instability/DNA repair syndromes
 - Skin chromosome breakage study (if not already performed)
 - The guidelines highlight the importance of genetic testing to identify family members who carry the mutation and are at risk for additional health consequences, such as a predisposition to cancer.
- Dyskeratosis congenita
 - Guidelines for diagnosis and management of DC were published by expert authors in consultation with a medical advisory board in 2015:
 - “Individuals should be considered to have DC if they have telomere lengths below the first percentile for age by multicolor flow cytometry with fluorescent in situ hybridization (flow FISH) in several subsets of leukocytes (granulocytes, CD45+ naïve T cells, CD45- memory T cells, CD20+ B cells, CD57+ NK/NKT cells) or a damaging or deleterious mutation in a DC associated gene”²²
- Shwachman-Diamond syndrome



- Draft consensus guidelines for the diagnosis and treatment of SDS were published in 2011:
 - “The clinical diagnosis is established by (a) documenting evidence of characteristic exocrine pancreatic dysfunction and hematological abnormalities and (b) excluding known causes of exocrine pancreatic dysfunction and bone marrow failure. Attention should be given to ruling out cystic fibrosis (the most common cause of pancreatic insufficiency) with a sweat chloride test, Pearson disease (pancreatic insufficiency and cytopenia, marrow ring sideroblasts and vacuolated erythroid and myeloid precursors), cartilage hair hypoplasia (diarrhea and cytopenia, and metaphyseal chondrodysplasia, and more common in certain isolated populations such as the Amish), and other inherited bone marrow failure syndromes (such as dyskeratosis congenita).”²³
 - “As the clinical diagnosis of SDS is usually difficult and patients may present at a stage when no clinical pancreatic insufficiency is evident, it is advisable to test most or all suspected cases for mutations in the *SBDS* gene. It is noteworthy that about 10% of patients with clinical features of SDS do not have identifiable mutations, and that *de novo* *SBDS* mutations have been identified in some families.”²³

Criteria

This guideline applies to inherited bone marrow failure syndrome multi-gene panel testing, which are defined as assays that simultaneously test for more than one inherited bone marrow failure gene

- Genetic Counseling
 - Pre and post-test genetic counseling by an appropriate provider (as deemed by the Health Plan policy), AND
- Previous Genetic Testing:
 - No known IBMFS pathogenic variant in the family, or
 - If there is a known IBMFS pathogenic variant in the family, testing has been performed and is negative, and a diagnosis of IBMFS is still suspected, AND
- The member has or is suspected to have a condition that will benefit from information provided by the requested IBMFS gene testing based on at least one of the following:
 - The member meets all criteria in a test-specific guideline, if available, OR
 - The member displays clinical features of the condition for which testing is being requested, by at least one of the following, AND:



Seattle Children's[®]
HOSPITAL • RESEARCH • FOUNDATION

PLUGS[®]

Patient-centered Laboratory
Utilization Guidance Services

© 2012–2020 Seattle Children's Hospital. All rights reserved

June 2020

- Unexplained chronic cytopenia with or without associated congenital physical anomalies consistent with the condition
- sporadic aplastic anemia
- myelodysplastic syndrome
- lack of cytopenias but classic physical findings, cancer diagnosis, or family history
- Acquired etiologies have been considered and ruled out when possible (e.g., immune-mediated or viral), AND
- Predicted impact on health outcomes, including immediate impact on medical management based on the molecular results , AND
- The member does not have a known underlying cause for their symptoms (e.g. known genetic condition), AND
- Rendering laboratory is a qualified provider of service per the Health Plan policy.

Exclusions and Other Considerations:

- If clinical screening tests are indicative of a specific IBMFS, a smaller multi-gene panel that contains condition specific gene will be reimbursed (i.e. Fanconi Anemia gene panel).
- Deletion/duplication analysis is often performed concurrently on clinically available panels. If deletion/duplication analysis is not performed on initial panel testing or is not comprehensive (i.e. doesn't detect smaller exon deletions) and sequencing is negative, consider sending for deletion/duplication analysis.
- Alternative sample, such as DNA from a skin biopsy may need to be considered in a patient with MDS/AML and/or when there is concern for somatic reversion events.¹

References

1. West AH, Churpek JE. Old and new tools in the clinical diagnosis of inherited bone marrow failure syndromes. *Hematology Am Soc Hematol Educ Program*. 2017;2017(1):79-87.
2. Ghemlas I, Li H, Zlateska B, et al. Improving diagnostic precision, care and syndrome definitions using comprehensive next-generation sequencing for the inherited bone marrow failure syndromes. *J Med Genet*. 2015;52(9):575-584.
3. Keel SB, Scott A, Sanchez-Bonilla M, et al. Genetic features of myelodysplastic syndrome and aplastic anemia in pediatric and young adult patients. *Haematologica*. 2016;101(11):1343-1350.



Seattle Children's[®]
HOSPITAL • RESEARCH • FOUNDATION

PLUGS[®]

Patient-centered Laboratory
Utilization Guidance Services

© 2012–2020 Seattle Children's Hospital. All rights reserved

June 2020

4. Mehta PA, Tolar J. Fanconi Anemia. In: Adam MP, Ardinger HH, Pagon RA, et al., eds. *GeneReviews*((R)). Seattle (WA)1993.
5. Savage SA, Alter BP. Dyskeratosis congenita. *Hematol Oncol Clin North Am.* 2009;23(2):215-231.
6. Glousker G, Touzot F, Revy P, Tzfati Y, Savage SA. Unraveling the pathogenesis of Hoyeraal-Hreidarsson syndrome, a complex telomere biology disorder. *Br J Haematol.* 2015;170(4):457-471.
7. Bertuch AA. The molecular genetics of the telomere biology disorders. *RNA Biol.* 2016;13(8):696-706.
8. Myers KC, Bolyard AA, Otto B, et al. Variable clinical presentation of Shwachman-Diamond syndrome: update from the North American Shwachman-Diamond Syndrome Registry. *J Pediatr.* 2014;164(4):866-870.
9. Nelson AS, Myers KC. Diagnosis, Treatment, and Molecular Pathology of Shwachman-Diamond Syndrome. *Hematol Oncol Clin North Am.* 2018;32(4):687-700.
10. Clinton C, Gazda HT. Diamond-Blackfan Anemia. In: Adam MP, Ardinger HH, Pagon RA, et al., eds. *GeneReviews*((R)). Seattle (WA)1993.
11. Arbiv OA, Cuvelier G, Klaassen RJ, et al. Molecular analysis and genotype-phenotype correlation of Diamond-Blackfan anemia. *Clin Genet.* 2018;93(2):320-328.
12. Klein C. Genetic defects in severe congenital neutropenia: emerging insights into life and death of human neutrophil granulocytes. *Annu Rev Immunol.* 2011;29:399-413.
13. Geddis AE. Congenital amegakaryocytic thrombocytopenia. *Pediatr Blood Cancer.* 2011;57(2):199-203.
14. Dokal I, Vulliamy T. Inherited bone marrow failure syndromes. *Haematologica.* 2010;95(8):1236-1240.
15. Hsu AP, McReynolds LJ, Holland SM. GATA2 deficiency. *Curr Opin Allergy Clin Immunol.* 2015;15(1):104-109.
16. Hirabayashi S, Wlodarski MW, Kozyra E, Niemeyer CM. Heterogeneity of GATA2-related myeloid neoplasms. *Int J Hematol.* 2017;106(2):175-182.
17. Chen DH, Below JE, Shimamura A, et al. Ataxia-Pancytopenia Syndrome Is Caused by Missense Mutations in SAMD9L. *Am J Hum Genet.* 2016;98(6):1146-1158.
18. Davidsson J, Puschmann A, Tedgard U, Bryder D, Nilsson L, Cammenga J. SAMD9 and SAMD9L in inherited predisposition to ataxia, pancytopenia, and myeloid malignancies. *Leukemia.* 2018;32(5):1106-1115.
19. Churpek JE. Familial myelodysplastic syndrome/acute myeloid leukemia. *Best Pract Res Clin Haematol.* 2017;30(4):287-289.
20. Liew E, Owen C. Familial myelodysplastic syndromes: a review of the literature. *Haematologica.* 2011;96(10):1536-1542.
21. *Fanconi Anemia: Guidelines for Diagnosis and Management.* Fourth ed: Fanconi Anemia Research Fund; 2014.



22. *Dyskeratosis congenita and telomere biology disorders: Diagnosis and management guidelines*. First ed. New York, NY: Dyskeratosis Congenita Outreach, Inc.; 2015.
23. Dror Y, Donadieu J, Kogelmeier J, et al. Draft consensus guidelines for diagnosis and treatment of Shwachman-Diamond syndrome. *Ann N Y Acad Sci*. 2011;1242:40-55.



Seattle Children's[®]
HOSPITAL • RESEARCH • FOUNDATION

PLUGS[®]

Patient-centered Laboratory
Utilization Guidance Services

© 2012–2020 Seattle Children's Hospital. All rights reserved

June 2020